

# BLNK (Phospho Tyr84) Rabbit pAb

CatalogNo: AT7176

## •Main information

<b>Target</b>	BLNK Phospho Tyr84
<b>Reactivity</b>	Human,Mouse,Rat
<b>Applications</b>	WB,IHC,IF,ELISA
<b>MW (kDa)</b>	65kD (Calculated) 65kD (Observed)
<b>Host Species</b>	Rabbit
<b>Isotype</b>	IgG
<b>Modified</b>	Phospho

## •Recommended Dilutions

WB 1:500-1:2000
IHC 1:100-1:300
ELISA 1:10000
IF 1:50-200 Note: For IHC, we suggest antigen retrieval with TE buffer pH 9.0 (Cat#RH0011)

## •Detailed information

**Specificity** Phospho-BLNK (Y84) Polyclonal Antibody detects endogenous levels of BLNK protein only when phosphorylated at Y84.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):EMyVM

**Storage** -15°C to -25°C/1 year(Do not lower than -25°C, Ship by ice bag)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

**Clonality** Polyclonal

---

## •Target Information

**Gene name** BLNK

**Protein Name** B-cell linker protein

Organism	Gene ID	UniProt ID
Human	<a href="#">29760</a> ;	<a href="#">Q8WV28</a> ;
Mouse	<a href="#">17060</a> ;	<a href="#">Q9QUN3</a> ;
Rat	<a href="#">499356</a> ;	<a href="#">Q4KM52</a> ;


**Cellular Localization** Cytoplasm . Cell membrane . BCR activation results in the translocation to membrane fraction.

**Tissue specificity** Expressed in B-cell lineage and fibroblast cell lines (at protein level). Highest levels of expression in the spleen, with lower levels in the liver, kidney, pancreas, small intestines and colon.


**Function** Disease:Defects in BLNK are the cause of hypoglobulinemia and absent B-cells [MIM:604515]. This is a developmental blockage at the pro- to pre-B-cell transition.,Disease:In 6 of 34 childhood pre-B acute lymphoblastic leukemia (ALL) samples that were tested showed a complete loss or drastic reduction of BLNK expression.,Function:Functions as a central linker protein that bridges kinases associated with the B-cell receptor (BCR) with a multitude of signaling pathways, regulating biological outcomes of B-cell function and development. Plays a role in the activation of ERK/EPHB2, MAP kinase p38 and JNK. Modulates AP1 activation. Important for the activation of NF-kappa-B and NFAT. Plays an important role in BCR-mediated PLCG1 and PLCG2 activation and Ca(2+) mobilization and is required for trafficking of the BCR to late endosomes. However, does not seem to be required for pre-BCR-mediated activation of MAP kinase and phosphatidylinositol 3 (PI3) kinase signaling. May be required for the RAC1-JNK pathway. Plays a critical role in orchestrating the pro-B cell to pre-B cell transition (By similarity). Plays an important role in BCR-induced B-cell apoptosis.,online information:BLNK mutation db,PTM:Following BCR activation, phosphorylated on tyrosine residues by SYK and LYN. When phosphorylated, serves as a scaffold to assemble downstream targets of antigen activation, including PLCG1, VAV1, GRB2 and NCK1. Phosphorylation of Tyr-84, Tyr-178 and Tyr-189 facilitates PLCG1 binding. Phosphorylation of Tyr-96 facilitates BTK binding. Phosphorylation of Tyr-72 facilitates VAV1 and NCK1 binding. Phosphorylation is required for both Ca(2+) and MAPK signaling pathways.,similarity:Contains 1 SH2 domain.,subcellular location:BCR activation results in the translocation to membrane fraction.,subunit:Associates with PLCG1, VAV1 and NCK1 in a B-cell antigen receptor-dependent fashion. Interacts with VAV3, PLCG2 and GRB2. Interacts through its SH2 domain with CD79A.,tissue specificity:Expressed in B-cell lineage and fibroblast cell lines (at protein level). Highest levels of expression in the spleen, with lower levels in the liver, kidney, pancreas, small intestines and colon.,

---

## •Validation Data



Immunohistochemical analysis of paraffin-embedded human brain, using BLNK (Phospho-Tyr84) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from K562 cells treated with starved 24h, using BLNK (Phospho-Tyr84) Antibody. The lane on the right is blocked with the phospho peptide.

## •Contact information

Orders: [order@lamarck.cn](mailto:order@lamarck.cn)  
Support: [support@lamarck.cn](mailto:support@lamarck.cn)  
Telephone: 400-801-6722  
Website: <http://www.Lamarck.cn>  
Address: All rights reserved ©2025 Lamarck.



Please scan the QR code to access additional product information:  
**BLNK (Phospho Tyr84) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)